

PRODUCT INFORMATION

Mitis Salivarius Agar

Cat. No. M13-117

DESCRIPTION

Agar medium used for the selective isolation of *Streptococcus mitis*, *Streptococcus Salivarius*, and Enterococcus from heavily contaminated specimens.

FORMULA (g/L)

Enzymatic Digest of Casein	15.0 g	Peptic Digest of Animal Tissue	5.0 g
Sucrose	50.0 g	Dextrose	1.0 g
Dipotassium Phosphate	4.0 g	Trypan Blue	0.075 g
Crystal Violent	0.0008 g	Agar	15.0 g

Final pH: 7.0 ± 0.2 at 25 °C

*Grams per liter may be adjusted or formula supplemented to obtain desired performance.

PREPARATION

Mix 90 grams of the medium in one liter of purified water until evenly dispersed. Heat with repeated stirring to dissolve completely. Distribute and autoclave at 121°C for 15 minutes. Cool to 45-50°C and aseptically add 1 ml of a 1% Potassium Tellurite solution.

QUALITY CONTROL SPECIFICATIONS

1. The powder is homogeneous, free flowing and light bluish-beige.
2. Visually the prepared medium is clear to slightly hazy and dark royal blue.
3. Expected cultural response after 18-48 hours at 35°C after preparation with
4. 1% Potassium Tellurite Solution.

ORGANISM	RESULT
<i>Escherichia coli</i> ATCC 25922	Inhibited
<i>Streptococcus mitis</i> ATCC 49456	Growth, Blue
<i>Streptococcus mutans</i> ATCC 25175	Growth, Blue
<i>Streptococcus salivarius</i> ATCC 7073	Growth, Blue "gum drop" colonies



STORAGE

Store the sealed bottle containing the dehydrated medium at 2 to 30°C. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect it from moisture and light. The dehydrated medium should be discarded if it is not free flowing or if the color has changed from the original color.

Version 01 – Date 06/20/24